Chem.-Biol. Interactions, 68 (1988) 305–306
Elsevier Scientific Publishers Ireland Ltd.

CHEMICO-BIOLOGICAL INTERACTIONS

CONTENTS

VOLUME 68 (1988)

Features of microsomal and cytosolic glutathione conjugation of hexachlorobutadiene in rat liver
by A. Wallin, R.G. Gerdes, R. Morgenstern, T.W. Jones and K. Ormstad (Stockholm, Sweden, Armidale, Australia and Baltimore, MD) 1

Trypsin hydrolysis of X-phenyl hippurates
by C.D. Selassie, M. Chow and C. Hansch (Claremont, CA) 13

Covalent binding of 4-nitrobenzyl mercaptan *S*-sulfate to the sulfhydryl groups of hepatic cytosolic proteins and bovine serum albumin with mixed disulfide bond formation
by K. Miwa, H. Okuda, S. Yoshioka and T. Watabe (Tokyo, Japan) 27

Porphinatoiron-mediated oxidation of polycyclic aromatic hydrocarbons
by A. Gold, K. Jayaraj, R. Sangaiah and L.M. Ball (Chapel Hill, NC) 39

Non-enzyme glutathione conjugation of 2-nitroso-6-methyldipyrido[1,2-*a*:3′,2′-*d*]imidazole (NO-Glu-P-1) in vitro: *N*-hydroxy-sulfonamide, a new binding form of arylnitroso compounds and thiols
by A. Umemoto, S. Grivas, Z. Yamaizumi, S. Sato and T. Sugimura (Tokyo, Japan) 57

Quantitation of interaction between cytochrome *P*-450scc and adrenodoxin — analysis in the median UV-region by second derivative spectroscopy
by V.M. Shkumatov, G. Smettan, O. Ristau, H Rein, K. Ruckpaul, V.L. Chashchin and A.A. Akhrem (Minsk, U.S.S.R. and Berlin, G.D.R.) 71

Identification of *S*-(2,5-dihydroxyphenyl)-cysteine and *S*-(2,5-dihydroxyphenyl)-*N*-acetylcysteine as urinary metabolites of acetaminophen in the mouse. Evidence for *p*-benzoquinone as a reactive intermediate in acetaminophen metabolism
by G.A. Pascoe, C.J. Calleman and T.A. Baille (Seattle, WA) 85

Covalent binding of acetaminophen to mouse hemoglobin. Identification of major and minor adducts formed in vivo and implications for the nature of the arylating metabolites
by D.B. Axworthy, K.-J. Hoffmann, A.J. Streeter, C.J. Calleman, G.A. Pascoe and T.A. Baillie (Seattle, WA) 99

Depurination and imidazole ring-opening in nucleosides and DNA alkylated by styrene oxide
by P. Vodička and K. Hemminki (Helsinki, Finland) 117

The influence of fluoranthene on the metabolism and DNA binding of benzo[*a*]pyrene in vivo in mouse skin
by J.E. Rice, M.C. Defloria, C. Sensenhauser and E.J. Lavoie (Piscataway, NJ and Valhalla, NY) 127

Four sulfhydryl-modifying compounds cause different structural damage but similar functional damage in murine lymphocytes
by D.D. Duncan and D.A. Lawrence (Albany, NY) 137

Phosphodiester cleavage in apurinic dinucleotides
by P. Vodička and K. Hemminki (Helsinki, Finland) 153

Formation of a pyridinium derivative by reaction of 4-hydroxypentenal with glycine
by S. Napetschnig, E. Schauenstein and H. Esterbauer (Graz, Austria) 165

In vivo and in vitro effects of a new macrolide antibiotic roxithromycin on rat liver cytochrome *P*-450: comparison with troleandomycin and erythromycin
by M. Delaforge, E. Sartori and D. Mansuy (Paris, France) 179

Streptozotocin-induced diabetes modulates the metabolic activation of chemical carcinogens
by C. Ioannides, S.L. Bass, A.D. Ayrton, J. Trinick, R. Walker and P.R. Flatt (Guildford, U.K.) 189
Study on the mechanism of interference of 3,4,3′,4′-tetrachlorobiphenyl with the plasma retinol-binding proteins in rodents
by A. Brouwer, W.S. Blaner, A. Kukler and K.J. Van Den Berg (Rijswijk and Wageningen, The Netherlands and New York, NY) 203
Induction of cytosolic and microsomal epoxide hydrolases in mouse liver by peroxisome proliferators, with special emphasis on structural analogues of 2-ethylhexanoic acid
by B. Lundgren, J. Meijer, W. Birberg, A. Pilotti and J.W. DePierre (Stockholm, Sweden) 219
Interaction of lysinoalanine with the protein synthesizing apparatus
by B.J. Lifsey Jr., W.R. Farkas and J.P. Reyniers (Knoxville, TN and Kingsville, TX) 241
Liver DNA alkylation after a single carcinogenic dose of dimethylnitrosamine to newborn and adult CFW Swiss mice
by P. Coccia, M. Salmona, L. Diomede, L. Citti, L. Mariani and M. Romano (Milan and Pisa, Italy) 259
Reductive addition of glutathione to *p*-benzoquinone, 2-hydroxy-*p*-benzoquinone and *p*-benzoquinone epoxides. Effects of the hydroxy- and glutathionyl substituents on *p*-benzohydroquinone autoxidation
by A. Brunmark and E. Cadenas (Linköping, Sweden) 273
Subject Index Volume 68 (1988) 299
Author Index Volume 68 (1988) 303
Contents Volume 68 (1988) 305